

## **Secondhand Marijuana/Cannabis Smoke**

“Smoke is smoke. Both tobacco and marijuana smoke impair blood vessel function similarly. People should avoid both, and governments who are protecting people against secondhand smoke exposure should include marijuana in those rules.”

*-Matthew Springer, cardiovascular researcher and Associate Professor of Medicine,  
University of California, San Francisco*

### **Facts about secondhand marijuana smoke:**

- Marijuana smoke is created by burning components of plants in the genus Cannabis.
- The terms marijuana and cannabis are often used interchangeably when discussing the smoking and vaping of the cannabis plant and its components, and the secondhand smoke created by combusting the product.
- Secondhand marijuana smoke is a complex chemical mixture of smoke emitted from combusted marijuana and the smoke that is exhaled by the user.
- Secondhand marijuana smoke contains fine particulate matter that can be breathed deeply into the lungs.
- Secondhand marijuana smoke contains many of the same cancer-causing substances and toxic chemicals as secondhand tobacco smoke. Some of the known carcinogens or toxins present in marijuana smoke include: acetaldehyde, ammonia, arsenic, benzene, cadmium, chromium, formaldehyde, hydrogen cyanide, isoprene, lead, mercury, nickel, and quinoline.<sup>i</sup>
- Marijuana smoke contains tetrahydrocannabinol (THC), the active chemical in cannabis.

### **Understanding the harms:**

Decades ago, people believed secondhand tobacco smoke presented no health risks to nonsmokers. Scientific research since that time changed this perception and led to smokefree environments. Likewise, scientific research is demonstrating that breathing secondhand marijuana smoke is dangerous to health and therefore should not be allowed in indoor spaces or places where people gather.

However, U.S. adults erroneously believe that exposure to cannabis secondhand smoke and cannabis smoking are safer than exposure to tobacco secondhand smoke and tobacco smoking, and these beliefs increased from 2017 to 2021.<sup>ii</sup> Likewise, cannabis is often seen as a harmless drug in the public eye, despite the fact that unequivocal evidence has established that cannabis smoking is harmful to the respiratory system, therefore the public should be educated about its potential harm.<sup>iii, iv</sup>

- As countries adopt more lenient cannabis use policies, perception of harm from secondhand cannabis smoke exposure is decreasing and most cannabis smoking is taking place at home. Research found that the perceived harm from secondhand cannabis smoke has declined as cannabis laws have expanded, and increasing the awareness of harms might reduce secondhand cannabis smoke exposure.<sup>v</sup>

## Health risks of exposure to secondhand marijuana smoke:

The 2024 U.S. Surgeon General's Report found that "secondhand cannabis smoke contains many of the same toxic and cancer-causing chemicals found in tobacco smoke, and some of those chemicals are found in higher amounts in secondhand cannabis smoke than in secondhand tobacco smoke."<sup>vi</sup>

Peer-reviewed and published studies indicate that exposure to secondhand marijuana smoke may have health and safety risks for the public, especially due to its similar composition to secondhand tobacco smoke.

- Secondhand smoke from combusted marijuana contains fine particulate matter that can be breathed deeply into the lungs,<sup>vii</sup> which can cause lung irritation, asthma attacks, and makes respiratory infections more likely. Exposure to fine particulate matter can exacerbate health problems especially for people with respiratory conditions like asthma, bronchitis, or COPD.<sup>viii</sup>
- Particulate levels from secondhand marijuana smoke are even higher than particulate levels from secondhand tobacco smoke. A study comparing indoor particulate matter 2.5 (PM2.5) levels from secondhand marijuana smoke and secondhand tobacco smoke concluded that "the average PM2.5 emission rate of the pre-rolled marijuana joints was found to be 3.5 times the average emission rate of Marlboro tobacco cigarettes, the most popular US cigarette brand."  
**Smoking a marijuana joint indoors can produce extremely high indoor PM2.5 concentrations**, thereby exposing the public and workers to dangerous secondhand marijuana smoke emissions.<sup>ix</sup>
- Being near people who are using inhaled cannabis is hazardous to human health. In a dispensary that allowed marijuana/cannabis smoking, research scientists discovered that the average PM2.5 emissions was 840 ug/m<sup>3</sup> over 9 visits, which exposed patrons and workers to air pollutant concentrations that are beyond hazardous levels.<sup>x</sup>
- The levels of PM2.5 fine particulate matter inside a Los Angeles cannabis consumption lounge were **10 times higher** during the lounge's busiest hours than the particulate levels inside the lounge when the business was closed and were 3 times higher than the levels during slower times of day.<sup>xi</sup>
- Ultrafine particles go deep into the lungs and can cause asthma, upper respiratory disease and more. Researchers studied the smoke from cannabis cigarettes and discovered that smoking cannabis in ozone-rich environments, both indoors and outdoors, will likely lead to ultrafine particle (UFP) formation.<sup>xii</sup>
- On-site consumption of cannabis using electric vaporizers, vape pens, and dab rigs produces a chemical aerosol that pollutes indoor air to unhealthy levels. This diminished air quality was observed when marijuana was not being combusted on-site. Researchers measured high levels of PM2.5 inside a marijuana retailer that allowed vaporizing, dabbing, and vaping marijuana - but did not allow smoking of marijuana or tobacco.<sup>xiii</sup>
- Researchers at the University of California at Berkeley studying fine particulate matter exposure found that cannabis bong smoking in the home generated **4 times greater PM2.5 concentrations** than cigarette or tobacco hookah (waterpipe) smoking.<sup>xiv</sup>
- Among adults living in a state with legalized medical cannabis and limited smokefree protections, researchers found that secondhand cannabis smoke exposure is common and associated with more frequent respiratory symptoms, particularly among cannabis users. Those exposed were more socially and economically vulnerable.<sup>xv</sup>

- In the first “whole-house” study of its kind, researchers measured both tobacco and marijuana secondhand smoke and found that PM2.5 concentrations from marijuana secondhand smoke were 4.4 times higher than emissions from secondhand tobacco smoke. Secondhand marijuana levels were higher than tobacco levels in every room of the house.<sup>xvi</sup>
- Children who live in households where cannabis is smoked indoors have biomarkers of cannabis secondhand smoke exposure that were 5 times higher than children who are not exposed to cannabis secondhand smoke in the home.<sup>xvii</sup>
- Significant amounts of mercury, cadmium, nickel, lead, hydrogen cyanide, and chromium, as well as 3 times the amount of ammonia, are found in mainstream marijuana smoke than in tobacco smoke.<sup>xviii</sup>
- In 2009, the California Office of Environmental Health Hazard Assessment added marijuana smoke to its Proposition 65 list of carcinogens and reproductive toxins, also known as the Safe Drinking Water and Toxic Enforcement Act of 1986. It reported that at least 33 individual constituents present in both marijuana smoke and tobacco smoke are Proposition 65 carcinogens.<sup>xix, xx</sup>
- Secondhand smoke from marijuana has many of the same chemicals as smoke from tobacco, including those linked to lung cancer.<sup>xxi</sup>
- Secondhand marijuana exposure impairs blood vessel function. Published studies on rats show that thirty minutes of exposure to secondhand marijuana smoke at levels comparable to those found in restaurants that allow cigarette smoking led to substantial impairment of blood vessel function. Marijuana smoke exposure had a greater and longer-lasting effect on blood vessel function than exposure to secondhand tobacco smoke.<sup>xxii</sup>
- One minute of exposure to marijuana SHS substantially impairs endothelial function in rats for at least 90 minutes, considerably longer than comparable impairment by tobacco SHS. The findings in rats suggest that SHS can exert similar adverse cardiovascular effects regardless of whether it is from tobacco or marijuana.<sup>xxiii</sup>
- Secondhand marijuana smoke and secondhand tobacco smoke is similar in many ways. More research is needed, but the current body of science shows that both tobacco and marijuana smoke have similar chemical composition and suggests that they may have harmful cardiovascular health effects, such as atherosclerosis (partially blocked arteries), heart attack, and stroke.<sup>xxiv</sup>
- Particle concentrations from dabbing and vaporizing cannabis can create levels of indoor air pollution similar as those seen in extreme air pollution events like wildfires and severe industrial pollution. Exposure at these concentrations can cause cardiovascular and respiratory disease.<sup>xxv</sup>
- Vaporized CBD produces significantly higher concentrations of formaldehyde, acetaldehyde, acrolein, diacetyl, and methylglyoxal compared with the other cannabinoid samples. Compared with other modalities of use for CBD and other cannabinoids, vaping has the potential to adversely impact human health by producing harmful products during the heated aerosolization process.<sup>xxvi</sup>
- People who are exposed to secondhand marijuana smoke can have detectable levels of THC (tetrahydrocannabinol) in their blood and urine.<sup>xxvii</sup>
- Marijuana also can be contaminated with mold, insecticides or other chemicals that may be released in secondhand smoke.<sup>xxviii</sup>

## Including Marijuana Smoking in Smokefree Public Place and Workplace Laws:

- Everyone has the right to breathe smokefree air. Smokefree policies are designed to protect the public and all workers from exposure to the health hazards caused by exposure to secondhand tobacco smoke. The same should be true for secondhand marijuana smoke.
- The 2024 U.S. Surgeon General's Report stated, "As states and communities consider whether use of cannabis in public places should be permitted, protecting all populations from exposure to secondhand tobacco smoke, including exposure to cannabis smoke, in all indoor settings should remain a priority."<sup>xxix</sup>
- The percent of U.S. adults who use marijuana more than doubled from 4.1% to 9.5% between 2001-2002 and 2012-2013,<sup>xxx</sup> which may also indicate an increase in exposure to secondhand marijuana smoke.
- The American Society for Heating, Refrigeration, and Air Conditioning Engineering (ASHRAE) is the organization that develops engineering standards for building ventilation systems. ASHRAE now bases its ventilation standard for acceptable indoor air quality on an environment that is completely free from secondhand tobacco smoke, secondhand marijuana smoke, and emissions from electronic smoking devices.<sup>xxxi</sup>
- To protect public health, improve consistency, and aid enforcement, smokefree laws for public places and workplaces should include tobacco as well as marijuana, whether it is smoked or aerosolized. Allowing marijuana smoking in places where smoking is now prohibited could undermine laws that protect the public from exposure to secondhand smoke. The Tobacco Control Legal Consortium issued an informative brief on [Lessons from Tobacco Control for Marijuana Regulation](#).<sup>xxxii</sup>
- Smokefree policies provide incentives to quit smoking, help denormalize smoking behavior, and are particularly effective among youth and young adults who are vulnerable to visual cues and social norms of smoking. It is likely that smokefree policies for marijuana will have a similar effect.
- As of April 1, 2025, 1095 localities and 40 states/territories/commonwealths restrict marijuana use in some or all smokefree spaces. Of these, 601 localities and 23 states/territories/commonwealths prohibit smoking and vaping of recreational and medical marijuana in one or more of the following venues: non-hospitality workplaces, restaurants, bars, and/or gambling facilities.

## Maintaining the Smokefree Gain from Industry Interference:

- The tobacco industry is setting its sights on the commercialization of marijuana and pushing for on-site consumption to once again normalize smoking and vaping in public, undermining decades of smokefree protections. Creating transparency about the relationships between the cannabis industry, related industries, and policymakers is essential to ensure appropriate regulation of cannabis products.
- An observational study found that nearly **half of lobbyists did not disclose that they represented the cannabis industry or interests**, which makes it difficult for people to track the industry's activity and spending. As the study notes, "The cannabis industry has an interest in creating a regulatory environment which maximizes profits at the cost of public health, similar to the tobacco, alcohol, and food industries." It is imperative for public health advocates to educate policymakers and the public about the cannabis industry's well-funded lobbying activities that aim to promote their profits over the public's wellbeing.<sup>xxxiii</sup>

- The 2024 U.S. Surgeon General's Report emphasized that "The increased legalization of recreational cannabis at state and local levels represents a threat to smokefree norms and protections and has implications for health disparities, due to creating exemptions for cannabis smoking" and advised that "smokefree policies will be most protective if they cover all emissions (i.e., smoke, aerosol) from the full range of products, including e-cigarettes, cigars, waterpipe, and cannabis."<sup>xxxiv</sup>

**In the interest of public health, the use of combustible or aerosolized marijuana should be prohibited wherever tobacco smoking is prohibited.**

## **ANR Foundation's Position on Exposure to Secondhand Marijuana Smoke:**

Marijuana smoke is a form of indoor air pollution. Therefore, ANR, our lobbying organization, includes marijuana within the definition of smoking, and all our model laws and policies include a prohibition on smoking or vaping marijuana wherever smoking and vaping is not allowed. Our organization does not have a position on whether marijuana should be legalized; we are committed to smokefree protections from secondhand smoke from tobacco products, marijuana and aerosol from electronic smoking devices.

**Nobody should have to breathe secondhand marijuana smoke at work, in public, or where they live.** We want healthy, smokefree air for workers and the public, so products like marijuana and electronic smoking devices (which can be used to vape a wide range of substances, including marijuana and hash oil) must not be used in smokefree environments where others are forced to breathe the secondhand emissions.

## **References**

- 
- <sup>i</sup> Moir, D., et al., A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chem Res Toxicol* 21: 494-502. (2008).  
<http://www.ncbi.nlm.nih.gov/pubmed/18062674>
- <sup>ii</sup> Janssen, F., et al., "Comparison between smoked tobacco and medical cannabis cigarettes concerning particulate matter," *Cannabis and Cannabinoid Research* [Online ahead of print], January 31, 2024.  
<https://www.doi.org/10.1089/can.2023.0201>
- <sup>iii</sup> Chambers, J., et al, "Perceptions of safety of daily cannabis vs tobacco smoking and secondhand smoke exposure, 2017-2021," *JAMA Network Open* 6(8): e2328691, August 11, 2023.  
<https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2808206>
- <sup>iv</sup> Khoj, L., et al., "Effects of cannabis smoking on the respiratory system: a state-of-the-art review," *Respiratory Medicine* 221: 107494, January 2024. <https://www.doi.org/10.1016/j.rmed.2023.107494>
- <sup>v</sup> Tripathi, O.; et.al., "Clearing the air: heightened perception of harm from secondhand cannabis smoke exposure is associated with no in-home cannabis smoking in a 21-country convenience sample," *Preventive Medicine* 189: 108178, December 2024. <https://doi.org/10.1016/j.ypmed.2024.108178>  
<https://www.sciencedirect.com/science/article/abs/S0091743524003335?via%3Dihub>
- <sup>vi</sup> U.S. Department of Health and Human Services. "Eliminating Tobacco-Related Disease and Death: Addressing Disparities," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, November 2024. <https://www.hhs.gov/surgeongeneral/reports-and-publications/tobacco/index.html>
- <sup>vii</sup> Hillier, FC.; et al. "Concentration and particle size distribution in smoke from marijuana cigarettes with different Δ9-tetrahydrocannabinol content." *Fundamental and Applied Toxicology*. Volume 4, Issue 3, Part 1, June 1984, Pages 451-454. <http://www.sciencedirect.com/science/article/pii/0272059084902021>
- <sup>viii</sup> "Air and Health: Particulate Matter." National Environmental Public Health Tracking Network, U. S. Environmental Protection Agency. <http://ephtracking.cdc.gov/showAirHealth.action#ParticulateMatter>
- <sup>ix</sup> Ott, W., et al., Measuring indoor fine particle concentrations, emission rates, and decay rates from cannabis use in a residence, *Atmospheric Environment: X*, Volume 10, 2021, 100106, ISSN 2590-1621,

---

<https://doi.org/10.1016/j.aeaoa.2021.100106>.

(<https://www.sciencedirect.com/science/article/pii/S259016212100006X>)

<sup>x</sup> PM2.5 Concentrations in the Smoking Lounge of a Cannabis Store, Huang, Murphy, Jacob and Schick, Environ. Sci. Technol. Lett. 2022, 9, 6, 551–556, Publication Date: May 26, 2022,

<https://doi.org/10.1021/acs.estlett.2c00148>, Copyright © 2022 American Chemical Society

<sup>xi</sup> Niu, M., et al., “Fine and ultrafine particle concentrations in a cannabis consumption lounge,” *Environmental Science and Technology Letters* [Online ahead of print], January 14, 2025.

<https://pubs.acs.org/doi/10.1021/acs.estlett.4c01019>

<sup>xii</sup> Yeh, K., et al., “Ultrafine particle generation from ozone oxidation of cannabis smoke,” *Environmental Science and Technology* 58(52): 22843-23252 December 31, 2024. <https://pubs.acs.org/doi/pdf/10.1021/acs.est.4c08311>

<sup>xiii</sup> Murphy, M.B.; Huang, A.S.; Schick, S.F., “PM2.5 concentrations in a cannabis store with on-site consumption,” *Environmental Health Perspectives* 129(6), June 16, 2021. <https://doi.org/10.1289/EHP8689>

<sup>xiv</sup> Nguyen., et al., Fine Particulate Matter Exposure From Secondhand Cannabis Bong Smoking (2022) [doi:10.1001/jamanetworkopen.2022.4744](https://doi.org/10.1001/jamanetworkopen.2022.4744).

<sup>xv</sup> Cohn, A.M., et al., “Secondhand cannabis smoke exposure and respiratory symptoms among adults living in a state with legalized medical cannabis with limited smoke-free protections,” *Preventive Medicine Reports* [Online ahead of print], July 23, 2024. <https://doi.org/10.1016/j.pmedr.2024.102835>

<sup>xvi</sup> Ott, W.R., et al., “Measuring PM2.5 concentrations from secondhand tobacco vs. marijuana smoke in 9 rooms of a detached 2-story house,” *Science of the Total Environment* 852 (2022) 158244.

<sup>xvii</sup> Tripathi, O., et.al., “Exposure to secondhand cannabis smoke among children,” *JAMA Network Open* 8(1): e2455963, January 23, 2025. <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2829482>

<sup>xviii</sup> Moir, D., et al., A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chem Res Toxicol* 21: 494-502. (2008).

<http://www.ncbi.nlm.nih.gov/pubmed/18062674>

<sup>xix</sup> “Evidence on the Carcinogenicity of Marijuana Smoke.” Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. August 2009.

[http://oehha.ca.gov/prop65/hazard\\_ident/pdf\\_zip/FinalMJsSmokeHID.pdf](http://oehha.ca.gov/prop65/hazard_ident/pdf_zip/FinalMJsSmokeHID.pdf)

<sup>xx</sup> Wang, X., et al., “Brief exposure to marijuana secondhand smoke impairs vascular endothelial function” (conference abstract). *Circulation* 2014; 130: A19538.

[http://circ.ahajournals.org/content/130/Suppl\\_2/A19538.abstract](http://circ.ahajournals.org/content/130/Suppl_2/A19538.abstract)

<sup>xxi</sup> “Evidence on the Carcinogenicity of Marijuana Smoke.” Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. August 2009.

[http://oehha.ca.gov/prop65/hazard\\_ident/pdf\\_zip/FinalMJsSmokeHID.pdf](http://oehha.ca.gov/prop65/hazard_ident/pdf_zip/FinalMJsSmokeHID.pdf)

<sup>xxii</sup> Wang, X., et al., “Brief exposure to marijuana secondhand smoke impairs vascular endothelial function” (conference abstract). *Circulation* 2014; 130: A19538.

[http://circ.ahajournals.org/content/130/Suppl\\_2/A19538.abstract](http://circ.ahajournals.org/content/130/Suppl_2/A19538.abstract)

<sup>xxiii</sup> Wang, X.; Derakhshandeh, R.; Liu, J.; Narayan, S.; Nabavizadeh, P.; Le, S.; Danforth, O.M.; Pinnamaneni, K.; Rodriguez, H.J.; Luu, E.; Sievers, R.E.; Schick, S.F.; Glantz, S.A.; Springer, M.L., “One minute of marijuana secondhand smoke exposure substantially impairs vascular endothelial function,” *Journal of the American Heart Association* 5(8): e003858, August 2016. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5015303/>

<sup>xxiv</sup> Springer, M.L.; Glantz, S.A. “Marijuana Use and Heart Disease: Potential Effects of Public Exposure to Smoke,” University of California at San Francisco. April 13, 2015.

<https://tobacco.ucsf.edu/sites/g/files/tkssra4661/f/u9/MSHS%20fact%20sheet%20for%20CA%204-13-15.pdf>

<sup>xxv</sup> Jaques, P., et al., “Measuring Aerosol Particle Emissions from Cannabis Vaporization and Dabbing”, Proceedings of the 15th Meeting of the International Society for Indoor Air Quality and Climate. July 22-27, 2018. Philadelphia, PA.

<sup>xxvi</sup> Robertson, N.E., et al., “Chemical composition of aerosols from the e-cigarette vaping of natural and synthetic cannabinoids,” *Chemical Research in Toxicology* 37(12): 1965-1975, December 16, 2024.

<https://www.doi.org/10.1021/acs.chemrestox.4c00326>

<sup>xxvii</sup> Herrmann ES, et al., “Non-smoker exposure to secondhand cannabis smoke II: Effect of room ventilation on the physiological, subjective, and behavioral/cognitive effects.” *Drug and Alcohol Dependence*. 2015 Jun 1;151:194-202. <http://www.ncbi.nlm.nih.gov/pubmed/25957157>

<sup>xxviii</sup> UC Davis Researchers, research letter published online in the journal of Clinical Microbiology and Infection, “A microbiome assessment of medical marijuana” April 2017 Volume 23, Issue 4, Pages 269–270

[https://www.clinicalmicrobiologyandinfection.com/article/S1198-743X\(16\)30605-X/pdf](https://www.clinicalmicrobiologyandinfection.com/article/S1198-743X(16)30605-X/pdf)

<sup>xxix</sup> U.S. Department of Health and Human Services. “Eliminating Tobacco-Related Disease and Death: Addressing Disparities,” U.S. Department of Health and Human Services, Centers for Disease Control and

---

Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, November 2024. <https://www.hhs.gov/surgeongeneral/reports-and-publications/tobacco/index.html>

<sup>xxx</sup> Hasin DS, et al. "Prevalence of Marijuana Use Disorders in the United States Between 2001-2002 and 2012-2013." JAMA Psychiatry. Published online October 21, 2015. <http://archpsyc.jamanetwork.com/article.aspx?articleid=2464591>

<sup>xxxii</sup> ANSI/ASHRAE Standard 62.1-2016 – Ventilation for Acceptable Indoor Air Quality. Atlanta, GA: American Society of Heating, Refrigerating, and Air-Conditioning Engineers, Inc. <https://www.ashrae.org/technical-resources/standards-and-guidelines/read-only-versions-of-ashrae-standards>

<sup>xxxiii</sup> Cork, Kerry. "Toking, Smoking & Public Health: Lessons from Tobacco Control for Marijuana Regulation." Tobacco Control Legal Consortium. June 2015. [http://www.publichealthlawcenter.org/sites/default/files/resources/tclc-synopsis-marijuana-tobacco-2015\\_0.pdf](http://www.publichealthlawcenter.org/sites/default/files/resources/tclc-synopsis-marijuana-tobacco-2015_0.pdf)

<sup>xxxiv</sup> U.S. Department of Health and Human Services. "Eliminating Tobacco-Related Disease and Death: Addressing Disparities," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, November 2024. <https://www.hhs.gov/surgeongeneral/reports-and-publications/tobacco/index.html>

<sup>xxxv</sup> Rotering, T., et al., "Cannabis industry lobbying in the Colorado state legislature in fiscal years 2010–2021," *International Journal of Drug Policy* 102: 103585, April 2022. <https://doi.org/10.1016/j.drugpo.2022.103585>

**May be reprinted with appropriate attribution to American Nonsmokers' Rights Foundation, © 2025**

*For more information:* [no-smoke.org/smokefree-threats/marijuana-smoke/](https://no-smoke.org/smokefree-threats/marijuana-smoke/)